

Literatur

1. Statistisches Bundesamt: Statistisches Jahrbuch 2014. URL www.destatis.de/DE/Publikationen/StatistischesJahrbuch/StatistischesJahrbuch.html Zugriff 11.01.15
2. Oberritter H, Schäbenthal K, von Ruesten A et al. (2013) The DGE-Nutrition Circle – Presentation and Basis of the Food-Related Recommendations from the German Nutrition Society (DGE). *Ernahrungs Umschau* 60: 24–29
3. Bergleiter S (2012) Nachhaltiger Fischkonsum: Ist die Empfehlung der DGE zum Fischverzehr unter Nachhaltigkeitsaspekten vertretbar? *Ernahrungs Umschau* 59: 282–285
4. Souci SW, Fachmann W, Kraut H. Die Zusammensetzung der Lebensmittel, Nährwert Tabellen. 7. Aufl., MedPharm Scientific Publishers, Stuttgart (2008)
5. Galli C, Rise P (2009) Fish consumption, omega 3 fatty acids and cardiovascular disease. *The science and the clinical trials. Nutr Health* 20: 11–20
6. König A, Bouzan C, Cohen JT et al. (2005) A quantitative analysis of fish consumption and coronary heart disease mortality. *Am J Prev Med* 29: 335–346
7. Noncommunicable diseases country profiles. WHO, Genf (2011)
8. Rapsomaniki E, Timmis A, George J et al. (2014) Blood pressure and incidence of twelve cardiovascular diseases: lifetime risks, healthy life-years lost, and age-specific associations in 1.25 million people. *Lancet* 383: 1899–1911
9. Dokholya RS, Albert CM, Appel LJ et al. (2004) A trial of omega-3 fatty acids for prevention of hypertension. *Am J Cardiol* 93: 1041–1043
10. Geleijnse JM, Giltay EJ, Grobbee DE et al. (2002) Blood pressure response to fish oil supplementation: meta-regression analysis of randomized trials. *J Hypertens* 20: 1493–1499
11. Mori TA, Burke V, Puddey IB et al. (2009) The effects of omega 3 fatty acids and coenzyme Q(10) on blood pressure and heart rate in chronic kidney disease: a randomized controlled trial. *J Hypertens* 27: 1863–1872
12. Ramel A, Alfredo MJ, Kiely M et al. (2010) Moderate consumption of fatty fish reduces diastolic blood pressure in overweight and obese European young adults during energy restriction. *Nutrition* 26:168–174
13. Theobald HE, Goodall AH, Sattar N et al. (2007) Low-dose docosahexaenoic acid lowers diastolic blood pressure in middle-aged men and women. *J Nutr* 137: 973–978
14. Hoshi T, Wissuwa B, Tian Y et al. (2013) Omega-3 fatty acids lower blood pressure by directly activating large-conductance Ca²⁺-dependent K(+)channels. *P Natl Acad Sci USA* 110: 4816–4821
15. Liu JC, Conklin SM, Manuck SB et al. (2011) Long-chain omega-3 fatty acids and blood pressure. *Am J Hypertens* 24: 1121–1126
16. Lara JJ, Economou M, Wallace AM et al. (2007) Benefits of salmon eating on traditional and novel vascular risk factors in young, non-obese healthy subjects. *Atherosclerosis* 193: 213–221
17. Schuchardt JP, Neubronner J, Kressel G et al. (2011) Moderate doses of EPA and DHA from re-esterified triacylglycerols but not from ethyl-esters lower fasting serum triacylglycerols in statin-treated dyslipidemic subjects: Results from a six month randomized controlled trial. *Prostaglandins Leukot Essent Fatty Acids* 85: 381–386
18. Dyerberg J, Madsen P, Moller JM et al. (2010) Bioavailability of marine n-3 fatty acid formulations. *Prostaglandins Leukot Essent Fatty Acids* 83: 137–141
19. Harris WS, De Caterina R, Marik PE (2013) Docosahexaenoic acid ethyl esters ineffective? *Proc Natl Acad Sci U S A* 110: E2259
20. Wu SY, Mayneris-Perxachs J, Lovegrove JA et al. (2014) Fish-oil supplementation alters numbers of circulating endothelial progenitor cells and microparticles independently of eNOS genotype. *Am J Clin Nutr* 100: 1232–1243
21. Cazzola R, Russo-Volpe S, Miles EA et al. (2007) Age- and dose-dependent effects of an eicosapentaenoic acid-rich oil on cardiovascular risk factors in healthy male subjects. *Atherosclerosis* 193: 159–167
22. Deutsche Gesellschaft für Ernährung e. V. (DGE) (2015) Evidenzbasierte Leitlinie: „Fettzufuhr und Prävention ausgewählter ernährungsmitbedingter Krankheiten“, 2. Version (2015)
23. Nelson RH (2013) Hyperlipidemia as a risk factor for cardiovascular disease. *Primary Care* 40:195–211
24. Browning LM, Krebs JD, Moore CS et al. (2007) The impact of long chain n-3 polyunsaturated fatty acid supplementation on inflammation, insulin sensitivity and CVD risk in a group of overweight women with an inflammatory phenotype. *Diabetes Obes Metab* 9: 70–80
25. Goodfellow J, Bellamy MF, Ramsey MW et al. (2000) Dietary supplementation with marine omega-3 fatty acids improve systemic large artery endothelial function in subjects with hypercholesterolemia. *J Am Coll Cardiol* 35: 265–270
26. Hartweg J, Perera R, Montori V et al. (2008) Omega-3 polyunsaturated fatty acids (PUFA) for type 2 diabetes mellitus. *Cochrane DB Syst Rev* 1: CD003205
27. Ciubotaru I, Lee YS, Wander RC (2003) Dietary fish oil decreases C-reactive protein, interleukin-6, and triacylglycerol to HDL-cholesterol ratio in postmenopausal women on HRT. *J Nutr Biochem* 14: 513–521
28. Skulas-Ray AC, Kris-Etherton PM, Harris WS et al. (2011) Dose-response effects of omega-3 fatty acids on triglycerides, inflammation, and endothelial function in healthy persons with moderate hypertriglyceridemia. *Am J Clin Nutr* 93: 243–252
29. Harris WS (1999) Nonpharmacologic treatment of hypertriglyceridemia: focus on fish oils. *Clin Cardiol* 22: II40–II43
30. Jump DB, Botolin D, Wan Y et al. (2008) Docosahexaenoic acid (DHA) and hepatic gene transcription. *Chem Phys Lipids* 153: 3–13
31. Grygiel-Gróniak B (2014) Peroxisome proliferator-activated receptors and their ligands: nutritional and clinical implications – a review. *Nutr J* 13: 17
32. Miller M, Langenberg P, Havas S (2007) Impact of lowering triglycerides on raising HDL-C in hypertriglyceridemic and non-hypertriglyceridemic subjects. *Int J Cardiol* 119: 192–195
33. Jacobson GA, Glickstein SB, Rowe JD et al. (2012) Effects of eicosapentaenoic acid and docosahexaenoic acid on low-density lipoprotein cholesterol and other lipids: a review. *J Clin Lipidol* 6: 5–18
34. Lankinen M, Kolehmainen M, Jaaskelainen T et al. (2014) Effects of whole grain, fish and bilberries on serum metabolic profile and lipid transfer protein activities: a randomized trial (Sysdimet). *Plos One* 9 (2)
35. Maki KC, van Elswyk ME, Alexander DD et al. (2012) A meta-analysis of randomized controlled trials that compare the lipid effects of beef versus poultry and/or fish con-

- sumption. *J Clin Lipidol* 6: 352–361
36. Adamsson V, Cederholm T, Vessby B et al. (2014) Influence of a healthy Nordic diet on serum fatty acid composition and associations with blood lipoproteins – results from the NORDIET study. *J Food Nutr Res* 58: 24114
 37. Bulliyya G (2002) Influence of fish consumption on the distribution of serum cholesterol in lipoprotein fractions: comparative study among fish-consuming and non-fish-consuming populations. *Asia Pac J Clin Nutr* 11: 104–111
 38. Calder PC (2006) n-3 polyunsaturated fatty acids, inflammation, and inflammatory diseases. *Am J Clin Nutr* 83: 1505S–1519S
 39. Caterina R, Madonna R, Bertolotto A (2007) n-3 fatty acids in the treatment of diabetic patients: biological rationale and clinical data. *Diabetes Care* 30: 1012–1026
 40. Li K, Huang T, Zheng J (2014) Effect of marine-derived n-3 polyunsaturated fatty acids on C-reactive protein, interleukin 6 and tumor necrosis factor α : a meta-analysis. *Plos One* 9: e88103
 41. Lopez-Garcia E, Schulze MB, Manson JE (2004) Consumption of (n-3) fatty acids is related to plasma biomarkers of inflammation and endothelial activation in women. *J Nutr* 134: 1806–1811
 42. Eschen O, Christensen JH, Caterina R et al. (2004) Soluble adhesion molecules in healthy subjects: a dose-response study using n-3 fatty acids. *Nutr Metab Cardiovas* 14: 180–185
 43. Hjerkin EM, Seljeflot I, Ellingsen I et al. (2005) Influence of long-term intervention with dietary counseling, long-chain n-3 fatty acid supplements, or both on circulating markers of endothelial activation in men with long-standing hyperlipidemia. *Am J Clin Nutr* 81: 583–589
 44. Miles EA, Thies F, Wallace FA (2001) Influence of age and dietary fish oil on plasma soluble adhesion molecule concentrations. *Clin Sci* 100:91–100
 45. Rees D, Miles EA, Banerjee T et al. (2006) Dose-related effects of eicosapentaenoic acid on innate immune function in healthy humans: a comparison of young and older men. *Am J Clin Nutr* 83: 331–342
 46. Bolton-Smith C, Woodward M, Tavendale R (1997) Evidence for age-related differences in the fatty acid composition of human adipose tissue, independent of diet. *Eur J Clin Nutr* 51: 619–624
 47. Jackson KG, Armah CK, Doman I et al. (2009) The impact of age on the post-prandial vascular response to a fish oil-enriched meal. *Brit J Nutr* 102: 1414–1419
 48. Geelen A, Brouwer IA, Schouten EG et al. (2004) Intake of n-3 fatty acids from fish does not lower serum concentrations of C-reactive protein in healthy subjects. *Eur J Clin Nutr* 58:1440–1442
 49. Madsen T, Christensen JH, Blom M et al. (2003) The effect of dietary n-3 fatty acids on serum concentrations of C-reactive protein: a dose-response study. *Brit J Nutr* 89: 517–522
 50. Vega-López S, Kaul N, Devaraj S et al. (2004) Supplementation with omega3 polyunsaturated fatty acids and all-rac alpha-tocopherol alone and in combination failed to exert an anti-inflammatory effect in human volunteers. *Metabolism* 53: 236–240
 51. Katan MB, Deslypere JP, vanBirgelen APJM et al. (1997) Kinetics of the incorporation of dietary fatty acids into serum cholesteryl esters, erythrocyte membranes, and adipose tissue: an 18-month controlled study. *J Lipid Res* 38: 2012–2022
 52. Patel PS, Bingham SA, Sharp SJ et al. (2009) Association between type of dietary fish and seafood intake and the risk of incident type 2 diabetes. The European Prospective Investigation of Cancer (EPIC)-Norfolk cohort study. *Diabetes Care* 32: 1857–1863
 53. Sotos PM, Guillen M, Sorli JV et al. (2011) Meat and fish consumption in a high cardiovascular risk spanish mediterranean population. *Nutri Hosp* 26: 1033–1040
 54. Djousse L, Gaziano JM, Buring JE et al. (2011) Dietary omega-3 fatty acids and fish consumption and risk of type 2 diabetes. *Am J Clin Nutr* 93: 143–150
 55. Ouellet V, Marois J, Weisnagel SJ et al. (2007) Dietary cod protein improves insulin sensitivity in insulin-resistant men and women. *Diabetes Care* 30: 2816–2821
 56. Huang T, Wahlqvist ML, Xu T et al. (2010) Increased plasma n-3 polyunsaturated fatty acid is associated with improved insulin sensitivity in type 2 diabetes in China. *Mol Nutr Food Res* 54: S112–S119
 57. Woodman RJ, Mori TA, Burke V et al. (2002) Effects of purified eicosapentaenoic and docosahexaenoic acids on glycemic control, blood pressure, and serum lipids in type 2 diabetic patients with treated hypertension. *Am J Clin Nutr* 76: 1007–1015
 58. Muley A, Muley P, Shah M (2014) ALA, fatty fish or marine n-3 fatty acids for preventing DM?: a systematic review and meta-analysis. *Curr Diabetes Rev* 10: 158–165
 59. Wallin A, Di Giuseppe D, Orsini N et al. (2012) Fish consumption, dietary long-chain n-3 fatty acids, and risk of type 2 diabetes. systematic review and meta-analysis of prospective studies. *Diabetes Care* 35: 918–929
 60. Zhang M, Picard-Deland E, Marette A (2013) Fish and marine omega-3 polyunsaturated fatty acid consumption and incidence of type 2 diabetes: a systematic review and meta-analysis. *Int J Endocrinol* 501015
 61. Zheng JS, Huang T, Yang J et al. (2012) Marine N-3 polyunsaturated fatty acids are inversely associated with risk of type 2 diabetes in Asians: a systematic review and meta-analysis. *Plos One* 7: e44525
 62. Chen R, Corona E, Sikora M et al. (2012) Type 2 diabetes risk alleles demonstrate extreme directional differentiation among human populations, compared to other diseases. *PLoS genetics* 8: e1002621
 63. Schulze MB, Hoffmann K, Manson JE et al. (2005) Dietary pattern, inflammation, and incidence of type 2 diabetes in women. *Am J Clin Nutr* 82: 675–684
 64. Wall R, Ross RP, Fitzgerald GF et al. (2010) Fatty acids from fish: the anti-inflammatory potential of long-chain omega-3 fatty acids. *Nutr Rev* 68: 280–289
 65. World Cancer Research Fund International (WCRF), American Institute for Cancer Research (AICR). Food, nutrition, physical activity, and the prevention of cancer: A global perspective. Washington DC, AICR (2007)
 66. Shen XY, Zhou JD, Dong JY (2012) Dietary intake of n-3 fatty acids and colorectal cancer risk: a meta-analysis of data from 489 000 individuals. *Br J Nutr* 108: 1550–1556
 67. Yu XF, Zou J, Dong J (2014) Fish consumption and risk of gastrointestinal cancers: A meta-analysis of cohort studies. *World J Gastroentero* 20: 15398–15412
 68. Liu J, Ma DWL (2014) The role of n-3 polyunsaturated fatty acids in the prevention and treatment of breast cancer. *Nutrients* 6: 5184–5223
 69. Patterson RE, Flatt SW, Newman VA (2011) Marine fatty acid intake is associated with

- breast cancer prognosis. *J Nutr* 141: 201–206
70. Zheng JS, Hu XJ, Zhao YM et al. (2013) Intake of fish and marine n-3 polyunsaturated fatty acids and risk of breast cancer: meta-analysis of data from 21 independent prospective cohort studies. *Brit Med J* 346
71. Saadatian-Elahi M, Norat T, Goudable J et al. (2004) Biomarkers of dietary fatty acid intake and the risk of breast cancer: a meta-analysis. *Int J Cancer* 111: 584–591
72. Schmidt JA, Gorst-Rasmussen A, Nystrom PW et al. (2014) Baseline patterns of adipose tissue fatty acids and long-term risk of breast cancer: a case-cohort study in the Danish cohort Diet, Cancer and Health. *Eur J Clin Nutr* 68: 1088–1094
73. Witt PM, Christensen JH, Schmidt EB et al. (2009) Marine n-3 polyunsaturated fatty acids in adipose tissue and breast cancer risk: a case-cohort study from Denmark. *Cancer Causes Control* 20: 1715–1721
74. Amin M, Jeyaganth S, Fahmy N et al. (2008) Dietary habits and prostate cancer detection: a case-control study. *Cuaj-Can Urol Assoc* 2: 510–515
75. Leitzmann MF, Stampfer MJ, Michaud DS et al. (2004) Dietary intake of n-3 and n-6 fatty acids and the risk of prostate cancer. *Am J Clin Nutr* 80: 204–216
76. Terry P, Lichtenstein P, Feychting M et al. (2001) Fatty fish consumption and risk of prostate cancer. *Lancet* 357: 1764–1766
77. Lovegrove C, Ahmed K, Challacombe B et al. (2015) Systematic review of prostate cancer risk and association with consumption of fish and fish-oils: analysis of 495,321 participants. *Int J Clin Pract* 69: 87–105
78. Pham TM, Fujino Y, Kubo T et al. (2009) Fish intake and the risk of fatal prostate cancer: findings from a cohort study in Japan. *Public Health Nutr* 12: 609–613
79. Chavarro JE, Stampfer MJ, Hall MN et al. (2008) A 22-y prospective study of fish intake in relation to prostate cancer incidence and mortality. *Am J Clin Nutr* 88: 1297–1303
80. Alexander D, Bassett J, Weed D et al. (2015) Meta-analysis of long-chain omega-3 polyunsaturated fatty acids (LCv-3PUFA) and prostate cancer. *Nutr Cancer* 67: 543–554
81. Chua ME, Sio MC, Sorongon MC et al. (2012) Relationship of dietary intake of omega-3 and omega-6 fatty acids with risk of prostate cancer development: a meta-analysis of prospective studies and review of literature. *Prostate Cancer* 2012: 826254
82. European Food Safety Authority (EFSA). Scientific opinion on the tolerable upper intake level of eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA) and docosapentaenoic acid (DPA). EFSA, (2012)
83. Nationale Verzehrsstudie II. Max Rubner Institut, Bundesforschungsinstitut für Ernährung und Lebensmittel, Karlsruhe (2008)
84. Lederman SA, Jones RL, Caldwell KL et al. (2008) Relation between cord blood mercury levels and early child development in a World Trade Center cohort. *Environ Health Persp* 116: 1085–1091
85. Mahaffey KR, Sunderland EM, Chan HM et al. (2011) Balancing the benefits of n-3 polyunsaturated fatty acids and the risks of methylmercury exposure from fish consumption. *Nutr Rev* 69: 493–508
86. Bundesinstitut für Risikobewertung (BfR). Aufnahme von Umweltkontaminanten über Lebensmittel. Ergebnisse des Forschungsprojektes LExUKon. URL www.bfr.bund.de/cm/350/aufnahme_von_umweltkontaminanten_ueber_lebensmittel.pdf Zugriff 11.01.15
87. European Food Safety Authority (EFSA) (2015) Statement on the benefits of fish/seafood consumption compared to the risks of methylmercury in fish/seafood. EFSA Journal 13: 3982
88. Bundesinstitut für Risikobewertung (BfR). Quecksilber und Methylquecksilber in Fischen und Fischprodukten – Bewertung durch die EFSA: Stellungnahme des BfR vom 29. März 2004. URL www.bfr.bund.de/cm/343/quecksilber_und_methylquecksilber_in_fischen_und_fischprodukten__bewertung_durch_die_efsa.pdf Zugriff 20.01.15

DOI: 10.4455/eu.2015.026